



 **Revuforj**[®]
(revumenib) tablets
25 mg • 110 mg • 160 mg

REVUFORJ READY

A doctor discussion guide for patients and caregivers throughout the treatment journey



All images are actor portrayals.

What is Revuforj?

Revuforj[®] (revumenib) is a prescription medicine used to treat adults and children 1 year and older with:

- acute leukemia with a lysine methyltransferase 2A gene (*KMT2A*) translocation whose disease has come back or has not improved after previous treatment(s)
- acute myeloid leukemia with a nucleophosmin 1 (*NPM1*) mutation whose disease has come back or has not improved after previous treatment(s) **and** who have no other satisfactory treatment options

Important Safety Information

What is the most important information I should know about Revuforj?

Revuforj may cause serious side effects, including:

Differentiation syndrome. Differentiation syndrome is a serious but common condition that affects your blood cells, which may be life-threatening or lead to death if not treated.

Please see additional Important Safety Information throughout and Full Prescribing Information, including **BOXED WARNINGS**, and Medication Guide and Instructions for Use.

MAKE THE MOST OF EVERY VISIT

It's important to partner with your healthcare team so you can learn what to expect while on treatment. Consider asking these questions so you can feel informed and supported while taking Revuforj® (revumenib).

Remember to take Revuforj exactly as your healthcare provider tells you to. Do not change your dose or stop taking Revuforj unless your healthcare provider tells you to.

Starting Revuforj

- When and how will I know if Revuforj is working?
- How long will I stay on Revuforj?
- Is Revuforj associated with any side effects I should be aware of?
- Should I avoid certain activities while taking Revuforj?
- Is it safe to eat particular foods or take other medications and/or supplements while taking Revuforj?
- How can I establish and stick to a treatment routine?

Support & Resources

- Where can I find support or advocacy groups for acute leukemia?
- Where can I find support or advocacy groups specifically for children and young adults?

Important Safety Information (*cont'd*)

What is the most important information I should know about Revuforj? (*cont'd*)
Differentiation syndrome (*cont'd*). Differentiation syndrome has happened as early as 3 days and up to 41 days after starting Revuforj. Tell any healthcare provider caring for you that you are taking a medicine that can cause differentiation syndrome. Call your healthcare provider or go to the nearest hospital emergency room right away if you develop any of the following symptoms of differentiation syndrome while taking Revuforj:

- | | | | |
|-----------------------|--------------------------------|---|-----------------------|
| • fever | • severe headache | • fast weight gain | • decreased urination |
| • cough | • confusion | • swelling of arms, legs, neck, groin, or underarm area | • rash |
| • shortness of breath | • dizziness or lightheadedness | | |

If you develop any of these symptoms of differentiation syndrome, your healthcare provider may start you on a medicine given through a vein (intravenous) called corticosteroids and may monitor you in the hospital.

Changes in electrical activity of your heart called QT prolongation. QT prolongation is a serious but common side effect that can cause irregular heartbeats that can be life-threatening, such as a heart attack, and can lead to death. Your healthcare provider will check the electrical activity of your heart with a test called an electrocardiogram (ECG) and will also do blood tests to check your potassium and magnesium levels before and during treatment with Revuforj. Tell your healthcare provider right away if you feel faint, lightheaded, dizzy, or if you feel your heart beating irregularly or fast during treatment with Revuforj.

Please see additional Important Safety Information throughout and Full Prescribing Information, including **BOXED WARNINGS, and Medication Guide and Instructions for Use.**

Here are some helpful tips to keep in mind when talking with members of your healthcare team:

- Before each appointment, think about any questions or information you or your loved one want to bring up with the doctor or the healthcare team
- Write your questions in a notebook so you don't forget to ask them—and so you can take notes during the visit
- Mention any symptoms or changes you've noticed since the last appointment
- Ask for written information about you or your loved one's test results and the treatment plan
- Don't be afraid to ask the doctor to slow down or explain things in another way if you don't understand
- Make sure you know who the main point of contact is, and who to contact after hours in case of an emergency

Notes

Important Safety Information (*cont'd*)

Before taking Revuforj® (revumenib), tell your healthcare provider about all your medical conditions, including if you:

- have any heart problems, including a condition called long QT syndrome
- have been told you have low blood levels of potassium or magnesium
- are pregnant or plan to become pregnant. Revuforj can harm your unborn baby

Females who are able to become pregnant

- Your healthcare provider will perform a pregnancy test within 7 days before you start treatment with Revuforj
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with Revuforj
- Use effective birth control (contraception) during treatment with Revuforj and for 4 months after the last dose of Revuforj

Males who have female partners who are able to become pregnant

- Use effective birth control during treatment with Revuforj and for 4 months after the last dose of Revuforj
- Talk to your healthcare provider about birth control methods you can use during this time
- Revuforj may cause fertility problems in females and males, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you
- are breastfeeding or plan to breastfeed. It is not known if Revuforj passes into your breast milk. Do not breastfeed during your treatment with Revuforj or for 1 week after your last dose of Revuforj

Please see additional Important Safety Information throughout and Full Prescribing Information, including **BOXED WARNINGS**, and Medication Guide and Instructions for Use.

 **Revuforj**[®]
(revumenib) tablets
25 mg • 110 mg • 160 mg

CAREGIVER CORNER

As a caregiver, you take on a lot of responsibility—and you may feel unsure about where to start. These discussion questions are geared toward your needs, so you can care for your loved one and yourself during their treatment journey.

- How can I better support my loved one while they're on treatment?
- Is there anything specific I should know about caring for someone living with acute leukemia?

In addition, **remember to make time for yourself** and take care of your own needs so that you stay healthy and well. Don't be afraid to reach out for help. You and your loved one are not alone.

Discover more tools and resources to help support you or your loved one at

[Revuforj.com](https://www.revuforj.com)

 **Revuforj**[®]
(revumenib) tablets
25 mg • 110 mg • 160 mg

Important Safety Information (cont'd)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Revuforj and other medicines may affect each other, causing side effects.

The most common side effects of Revuforj include:

- infections, including bacterial and viral
- changes in liver function tests
- bleeding (hemorrhage)
- nausea and vomiting
- muscle pain
- low white blood cell counts with fever
- diarrhea
- tiredness
- swelling in the arms and legs
- decreased appetite
- constipation

Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with Revuforj if you develop certain side effects. These are not all the possible side effects of Revuforj. Call your healthcare provider for medical advice about side effects.

You are encouraged to report side effects of prescription drugs to FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and Full Prescribing Information, including BOXED WARNINGS, and Medication Guide and Instructions for Use.

The information provided here is not a substitute for talking with your healthcare provider. Your healthcare provider is the best source of information about your disease.



Revuforj and the Syndax logos are registered trademarks of Syndax.
© 2025 Syndax Pharmaceuticals, Inc. REJ-US-250180 11/2025